



FIFRA
Section 24(c) Special Local Need Label

RESTRICTED USE PESTICIDE
Due to Toxicity in Fish, Mammals, and Aquatic Organisms

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ALABAMA

ZORO® Miticide/Insecticide

EPA Reg. No. 67760-71
EPA SLN No. AL -070005

For control of spider mites in cotton

Active Ingredients:*	
Abamectin.....	1.9%*
Other Ingredients:	98.1%
Total:	100.0%

*1 gal. Contains 0.15 lb. abamectin

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered ZORO Miticide/Insecticide label.

This label must be in the possession of the user at the time of pesticide application.

GENERAL INFORMATION

Use Restrictions

- Do not apply more than 32 fl. oz. or more than 2 applications of ZORO/A in any growing season.
- Do not apply within 21 days of last treatment.
- Do not apply with 20 days of harvest.
- Do not graze or feed cotton foliage.
- Do not apply this product through any type of irrigation system.

Note: Do not apply by ground within 25 ft., or by air within 150 ft. of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

CROP USE DIRECTIONS

Cotton

ZORO Miticide/Insecticide is an emulsifiable concentrate containing 0.15 lb. abamectin/gal. which, when mixed with water according to the Directions for Use, will control spider mites (two-spotted, and Carmine) on cotton.

For control of spider mites on cotton, mix with water and apply at the rates given in the following table:

Spider Mites	Fl. oz./A	
	Mid-season/Lay-by	Early Season
Two-spotted Carmine	8-16	4-6

Use 8-16 fl. oz. of ZORO/A depending on size of the plant and density of the foliage. Apply with ground sprayers or aerially by fixed-wing aircraft or helicopter in a minimum of 5 gals. of water/A to provide thorough coverage. Begin application when mites are first noticed and repeat if necessary. For use on early season cotton, apply when cotton is under 10 inches in height. Apply at a rate of 4-6 fl. oz. of ZORO/A by ground equipment only. Do not use less than 4 fl. oz./A.

Chemigation: **DO NOT** apply this product through any type of irrigation system.

Expiration Date:

24(c) registrant:
Cheminova, Inc.
One Park Drive, Suite 150
P.O. Box 110566
Research Triangle Park, NC 27709
www.cheminova.us.com

ZORO® is a registered trademark of Cheminova