



## REPORT™ HERBICIDE

TANK MIX WITH EVEREST HERBICIDE  
FOR WEED CONTROL IN WHEAT

EPA Reg. No. 67760-81

**GENERAL INFORMATION**

REPORT Herbicide is a dry flowable formulation that selectively controls certain broadleaf weeds in wheat. The degree and duration of weed control depends on the weed spectrum and infestation intensity, the weed size at the time of application, and/or the environmental conditions at the following treatment.

REPORT is noncorrosive to equipment, nonflammable, nonvolatile, and does not freeze.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REPORT may be tank mixed with EVEREST Herbicide for improved control of weeds in wheat. Read and follow all label instructions regarding timing of application, precautions, and warnings for these herbicides before using the tank mixture. Follow the most restrictive labeling.

**POSTEMERGENCE APPLICATION**

**Winter Wheat:** Apply in the fall or spring anytime after the crop has 2 total leaves on main stem and any number of tillers. Apply before jointing begins.

Treat late-seeded winter wheat after the crop has started to tiller since the combined effect of herbicide stress and stress from cold weather and/or moisture could cause crop injury.

**Spring Wheat:** Apply anytime after emergence, but before the majority of plants have 4 total leaves on the main stem plus 2 tillers. Do not apply after jointing begins. Do not apply to durum wheat.

The addition of 2,4-D or Dicamba to the REPORT + EVEREST tank-mix is required when applying to spring wheat.

The 2,4-D and Dicamba rates are as follows:

- 2,4-D Amine or LV Ester (4lbs./gallon): 0.25-0.75 pt./acre
- 2,4-D LV Ester (6lbs./gallon):0.17-0.5pt./acre
- Dicamba (4 lbs./gallon): 2-4 fl. oz. per acre

**Note:** When REPORT + EVEREST is applied in a tank-mix with a dicamba-containing herbicide, wild oat control may be reduced.

**ADDITIONAL TANK MIX INFORMATION AND PRECAUTIONS**

**Tank Mixes with Insecticides:** Under certain conditions (stress from drought, cold weather or warm days and cold nights following application, or crops in the 2-4 leaf stage), tank mixtures or sequential treatments of REPORT + EVEREST and organophosphate insecticides (such as methyl parathion, "Di-Syston", etc.) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area first. If no symptoms of crop injury occur 14 days after treatment, treat the rest of the acreage.

Do not use REPORT + EVEREST plus malathion, as crop injury may result.

Do not apply REPORT + EVEREST within 60 days of crop emergence where an organophosphate insecticide (such as "Di-Syston") has been applied as an in-furrow treatment, as crop injury may result.

## SURFACTANTS

### Winter Wheat

**Non-ionic Surfactant (NIS):** Unless otherwise specified, add a nonionic surfactant having at least 80% active ingredient at 0.5% v/v (2 quarts per 100 gallons of spray solution). Antifoaming agents may be used if needed.

**Ammonium Nitrogen Fertilizer:** In addition to a non-ionic surfactant, use 2 qt./acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2lbs./acre of a spray-grade ammonium sulfate (AMS). Use 4 qt./acre UAN or 4lbs./acre AMS under arid conditions.

Do not use low rates of liquid fertilizer as a substitute for surfactant.

### Carrier Solutions (With Liquid Nitrogen Solution Fertilizer):

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing REPORT and EVEREST in fertilizer solution.

REPORT and EVEREST must be first slurried with water, then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while REPORT and EVEREST are added. Note that use of fertilizer as the carrier may result in temporary crop yellowing and stunting.

If using rates of liquid nitrogen fertilizer in the spray solution that are less than 50% of the spray solution volume, the addition of non-ionic surfactant is necessary. Add surfactant at 0.25% v/v (1 quart per 100 gallons of spray solution) based on local recommendations.

When using rates of liquid nitrogen fertilizer in the spray solution that are greater than 50% of the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer or consultant for a specific recommendation before adding surfactant to these tank mixtures.

### Spring Wheat

**Non-ionic Surfactant (NIS):** Unless otherwise specified, add a nonionic surfactant having at least 80% active ingredient at 0.125% v/v (1 pint per 100 gallons of spray solution). Antifoaming agents may be used if needed. DO NOT add NIS if also tank-mixing with 2,4-D LV Ester.

## RATE OF APPLICATION AND WEEDS CONTROLLED OR PARTIALLY CONTROLLED

A tank mixture of REPORT + EVEREST will provide control, partial control, and/or suppression of the following weeds when applied at the specified rates and application timing.

### GRASS WEED RECOMMENDATIONS

Grass Weeds	Early* Applications of 0.6 oz "Everest" + 0.3 oz. REPORT per acre	Early* Applications of 0.4 oz "Everest" + 0.2 oz. REPORT per acre	Late* Applications of 0.6 oz "Everest" + 0.3 oz. REPORT per acre	Late* Applications of 0.4 oz "Everest" + 0.2 oz. REPORT per acre
Annual Ryegrass***	C**	PC	C	PC
Downy Brome ( <i>Bromus tectorum</i> )	PC	NR	PC	NR
Cheat ( <i>Brome secalinus</i> )	C	C#	C	C#
Japanese Brome ( <i>Bromus japonicus</i> )	C	C#	C	C#
Wild Oat	C	C	C	C

\*EARLY APPLICATION: Timing – 2 leaf stage to January 1

LATE APPLICATION: Timing-after January 1 but prior to joint

\*\*C=Control

PC=Partial Control (partially controlled weeds exhibit a visual reduction in numbers and/or a significant loss of vigor)

NR=Not recommended

\*\*\*If REPORT + EVEREST is applied in a tank-mix combination with a 2,4-D containing broadleaf herbicide. Ryegrass control may be reduced

# For control of these weed species at the reduced rate, liquid nitrogen fertilizer as a carrier solution of 50% or greater must be added to the tank mix.

### BROADLEAF WEED RECOMMENDATIONS

Broadleaf Weeds	Early* Application of 0.6 oz "Everest" + 0.3 oz. REPORT per acre	Early* Application of 0.4 oz "Everest" + 0.2 oz. REPORT per acre	Late* Application of 0.6 oz "Everest" + 0.3 oz. REPORT per acre	Late* Application of 0.4 oz "Everest" + 0.2 oz. REPORT per acre
Bedstraw	PC**	NR	PC	NR
Black Mustard	C	C	C	C
Blue Mustard	C	C	C	C
Bur beakchervil	C	NR	C	NR
Bushy Wallflower/ Treacle Mustard	C	C	C	C
Buttercup	C	NR	C	NR
Canada thistle	PC	NR	PC	NR
Coast Fiddleneck (tarweed)	C	NR	C	NR
Common chickweed	C	NR	C	NR
Common groundsel	C	NR	C	NR
Conical catchfly	C	C	C	C
Corn gromwell	PC	NR	PC	NR
Curly dock	C	C	C	C
Corn spurry	C	NR	C	NR
Cow cockle	C	NR	C	NR
Cutleaf eveningprimrose	C	C	C	C
False chamomile	C	NR	C	NR
Falseflax	C	NR	C	NR
Field pennycress	C	C	C	C
Flixweed	C	C	C	C
Hempnettle	C	C	C	C
Henbit	C	C	C	C
Kochia+	PC	NR	PC	NR
Ladysthumb	C	NR	C	NR
Lambsquarter	C	NR	C	NR
Mayweed	C	C	C	C
Miners lettuce	C	C	C	C
Mouseear chickweed	C	NR	C	NR
Pennsylvania smartweed	PC	NR	PC	NR
Pineappleweed	C	C	C	C
Pigweeds (redroot, smooth, prostrate, tumble)	C	C	C	C
Prickly lettuce+	PC	NR	PC	NR
Prostrate knotweed	PC	NR	PC	NR
Purslane (common)	C	NR	C	NR
Redstem filaree	C	NR	C	NR
Russian thistle	PC	NR	PC	NR
Shepherd's purse	C	C	C	C
Speedwell	PC	NR	PC	NR
Sunflower	PC	NR	PC	NR
Tansymustard	C	C	C	C
Volunteer canola	C	NR	C	PC

Waterpod	C	C	C	C
White cockle	C	NR	C	NR
Wild buckwheat	PC	NR	PC	NR
Wild carrot	C	NR	C	NR
Wild garlic/wild onion	PC	NR	PC	NR
Wild mustard	C	C	C	C
Wild radish	PC	NR	PC	NR
Wild turnip	C	NR	C	NR

\*EARLY APPLICATION: Timing – 2 leaf stage to January 1

LATE APPLICATION: Timing – after January 1 but prior to joint

\*\*C=Control

PC=Partial Control (partially controlled weeds exhibit a visual reduction in numbers and/or a significant loss of vigor)

NR=Not recommended

±Naturally occurring resistant biotypes of these weeds are known to occur. See Tank Mixtures and Resistance sections on the REPORT label for additional information.

Refer to the complete REPORT and EVEREST labels for additional information regarding use restrictions, tank-mix partners, rotational cropping recommendations, sprayer cleanup, use precautions, and other information. The most restrictive provisions on either label will apply. Do not use this tank mix if any restrictions on the EVEREST label conflict with recommendations on the REPORT label.

## IMPORTANT

### **BEFORE USING REPORT, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

**Read the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label.**

This labeling must be in the possession of the user at the time of pesticide application.

REPORT™ is a trademark of Cheminova

EVEREST is a registered trademark of Arvesta Corporation.

Cheminova Inc.  
One Park Drive, Suite 150  
P.O. Box 110566  
Research Triangle Park, NC 27709  
www.cheminova.us.com